

QUINCLORAC 1.5L SELECT[™]

Selective post-emergent herbicide for efficient weed control

ACTIVE INGREDIENT

Quinclorac – Group 4

% ACTIVE INGREDIENT

Quinclorac 18.92%

FORMULATION

Flowable concentrate

CHEMICAL FAMILY

Quinoline carboxylic acid

BRAND ALTERNATIVE

Drive[®] XLR8

MODE OF ACTION

Synthetic auxin

RESTRICTED USE

No

SIGNAL WORD


Caution

PACKAGING

4 x 0.5 gal, 12 x 7.5 oz

LABELED USES

Residential and non-residential turfgrass



Quinclorac 1.5L Select[™] is a selective, post-emergent herbicide that can be applied to many types of turf, both residential and non-residential, to control a variety of weeds. **Quinclorac 1.5L Select[™]** can be applied from 7 days before seeding and up to 28 days after emergence on selected weeds.

PRODUCT BENEFITS

- Provides up to 90 days of control
- Works on both broadleaf and grassy weeds
- Selective control means it doesn't harm turf
- Controls crabgrass
- Use conjunction with seeding/overseeding
- Add **Duo Stick Select[™]** to enhance efficacy



Prime Source, a Division of Albaugh, LLC

1525 NE 36th Street, Ankeny, IA 50021 • www.primesource-albaugh.com

©2021 Albaugh, LLC. Prime Source and Select are trademarks of Albaugh, LLC. Drive is a registered trademark of BASF. Always read and follow label instructions.

QUINCLORAC 1.5L SELECT™

Selective post-emergent herbicide for efficient weed control

GENERAL INFORMATION

Quinclorac 1.5L Select™ is an auxin agonist and is classified as a quinoline carboxylic acid. It is absorbed by foliage and roots and translocated throughout the plant. The control symptoms exhibited by broadleaf weeds include leaf and stem curl or twisting, and chlorosis. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death. Apply **Quinclorac 1.5L Select™** to actively growing weeds as post-emergent broadcast or spot sprays using the turf species, rate and growth stages indicated on the label.

The use of **Quinclorac 1.5L Select™** before or after seeding or overseeding a turf area will not significantly interfere with the turfgrass seed germination and growth of those grass types identified as tolerant or moderately tolerant on the label.

Broadleaf weeds controlled

Bindweed, field	Geranium, Carolina
Clover, hop	Medic, black
Clover, red	Morningglory spp.
Clover, white	Speedwell, common
Daisy, English ^{2,5}	Speedwell, slender
Dandelion, common ²	Speedwell, thymeleaf
Dollarweed	Violet, wild

Grassy weeds suppressed

Barnyardgrass	Foxtail, yellow ¹
Crabgrass, large ^{1,4}	Kikuyugrass ^{2,3}
Crabgrass, smooth ^{1,4}	Signalgrass, broadleaf
Foxtail, giant ¹	Torpedograss ³
Foxtail, green ¹	

¹ Under certain conditions, application of **Quinclorac 1.5L Select™** made to annual grasses at 2 to 4 tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of **Quinclorac 1.5L Select™** + methylated seed oil are applied either before second tiller or as weed grasses mature.

² Tank mix partner or sequential application required.

³ Make 2 sequential applications of 1.0 fl. oz. (0.51 lb. AI/A) of **Quinclorac 1.5L Select™** per 1,000 sq. ft. and an additional sequential application up to 0.90 fl. oz. (0.46 lb. AI/A) of **Quinclorac 1.5L Select™** per 1,000 sq. ft. at 14- to 21-day intervals.

⁴ Biotypes of large and smooth crabgrass in California have shown varied response to **Quinclorac 1.5L Select™**. If control failure occurs following a full or split application, **DO NOT** reapply **Quinclorac 1.5L Select™**. Change to a herbicide with a different mode of action.

⁵ **NOT FOR USE** in California.

Refer to label for full list of weeds controlled.