

# OXEM 4 ME SELECT™

Systemic, protective and curative fungicide for turf and ornamentals

## ACTIVE INGREDIENT

Metalaxyl-M  
(mefenoxam) – Group 4

## % ACTIVE INGREDIENT

45.3%

## FORMULATION

Microemulsion concentrate

## CHEMICAL FAMILY

Acylamine

## BRAND ALTERNATIVE

Subdue Maxx®

## MODE OF ACTION

Selective inhibition of ribosomal RNA synthesis by affecting the activity of the RNA polymerases.

## RESTRICTED USE

No

## SIGNAL WORD

Warning

## PACKAGING

4 x 0.5 gal

## LABELLED USES

Turfgrass

Ornamental plants

Ornamental bulb, corm,  
and tuber plants

Christmas trees and  
Christmas tree plantations

Non-bearing citrus trees  
and fruit and nut plants

Retail nurseries

Residential and commercial  
landscapes, and interior  
plantscape ornamentals



**Oxem 4 ME Select™** is a systemic turf and ornamental fungicide containing mefenoxam for the control of damping off, root and stem diseases caused by *Pythium* and *Phytophthora* spp., and foliar diseases such as downy mildew and those caused by *Phytophthora* spp. It works by suppressing essential components of the disease so that the infection doesn't grow or spread.

## PRODUCT BENEFITS

- Broad-spectrum control of *Pythium* and *Phytophthora* diseases
- Systemic activity provides root to leaf protection
- Provides protective and curative disease control
- Can be used on cool and warm season grasses
- 2X the concentration of competitive standard formulations
- Excellent tank-mix partner



**Prime Source, a Division of Albaugh, LLC**

1525 NE 36th Street, Ankeny, IA 50021 • [www.primesource-albaugh.com](http://www.primesource-albaugh.com)

©2022 Albaugh, LLC. Prime Source and the water drop logo are registered trademarks of Albaugh, LLC. Subdue Maxx is a registered trademark of a Syngenta Group Company. Always read and follow label Precautions and Directions For Use.

# OXEM 4 ME SELECT™

Systemic, protective and curative fungicide for turf and ornamentals

## Label use instructions for turf, flowers and woody ornamentals

APPLICATION AREA	DISEASES CONTROLLED
<b>TURF:</b> including but not limited to golf courses, lawns, landscape areas around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, and athletic fields.	Downy mildew <i>Pythium</i> blight <i>Pythium</i> damping-off
<b>BEDDING PLANTS:</b> <i>Ageratum</i> , Algerian ivy, <i>Artemisia</i> , <i>Aster</i> , begonia, <i>Caladium</i> , carnation, chrysanthemum, <i>Coleus</i> , daisy, English ivy, foxglove, <i>Gaillardia</i> , geranium, <i>Impatiens</i> , marigold, pansy, petunia, <i>Phlox</i> , pinks, primrose, prostrate rosemary, <i>Salvia</i> , snapdragon, <i>Verbena</i> , <i>Vinca</i> , zinnia, including all cultivars, varieties, and hybrids of these.	Damping-off, root diseases, and stem diseases caused by <i>Pythium</i> and <i>Phytophthora</i> spp.  Foliar diseases such as downy mildew and foliar diseases caused by <i>Phytophthora</i> spp., including <i>P. ramorum</i>
<b>FLOWERS</b> (including bulb, corm, and tuber plants): African violet, <i>Anthurium</i> , baby's breath, carnation, chrysanthemum, columbine, <i>Delphinium</i> , Easter lily, geranium, <i>Gloxinia</i> , poinsettia, rose, including all cultivars, varieties, and hybrids.	Downy mildew, <i>Phytophthora</i> spp., <i>Pythium</i> spp.
<b>FOLIAGE PLANTS:</b> <i>Aglaonema</i> , <i>Aphelandra</i> , <i>Dieffenbachia</i> , <i>Peperomia</i> , philodendron, pothos, <i>Schefflera</i> , <i>Sedum</i> , <i>Sempervivum</i> , <i>Zygocactus</i> , including all cultivars, varieties, and hybrids of these.	
<b>WOODY ORNAMENTALS:</b> <i>Aucuba japonica</i> , arborvitae, azaleas, boxwood, <i>Ceanothus</i> , <i>Cotoneaster</i> , dogwood, <i>Ficus</i> , "Halls" honeysuckle, <i>Ilex</i> , <i>Juniperus</i> spp., <i>Photinia</i> , <i>Pieris japonica</i> , <i>Pinus</i> spp., <i>Pittosporum</i> , <i>Rhododendron</i> , white cedar, white pine, yew, including all cultivars, varieties, and hybrids.	<i>Pythium</i> and <i>Phytophthora</i> root and crown rot  Downy mildew, <i>Phytophthora</i> spp., <i>Pythium</i> spp.
Conifers in nurseries and forest nurseries (including Christmas trees)	<i>Phytophthora</i> root and stem diseases, <i>Phytophthora</i> foliar diseases
Conifers in plantations (including Christmas trees)	<i>Phytophthora</i> root, stem, and foliar diseases  Foliar diseases caused by <i>Phytophthora</i> spp., including <i>P. ramorum</i>
<b>CITRUS</b> in nurseries and landscape plantings (non-bearing)	Citrus foot rot, root rot, trunk canker caused by <i>Phytophthora</i> spp.
Fruit and nut plants in nurseries and landscape plantings (non-bearing)	<i>Phytophthora</i> root, crown, and collar rot, <i>Pythium</i> root rot

See label for full list of diseases controlled and application rates.